

Non-Point Source Contamination and Poorly Sealed Wells in the Treasure Valley Aquifer

Shawn Benner

Hydrology Program

Boise State University

Idaho Water Resources Research Institute

Shallow Treasure Valley Aquifer






Fig.4

Arsenic is an example of a widespread, non-point source contaminant in the Treasure Valley




Monty Busbee, Graduate Student


Vertical Distribution of Arsenic is Also Heterogeneous



Monty Busbee, Graduate Student




**Contaminated waters are heterogeneous, some wells are clean, others are not. We are not yet able to predict
Where contamination will be elevated**




**New wells can change flow paths and
CHANGE what water goes to a distant well,
with a potential to induce contamination.**

Lake Lowell




Importantly, the way in which a poorly sealed well will induce a change in flow is not easily predicted



Importantly, the way in which a poorly sealed well will induce a change in flow is not easily predicted

Other Emerging Non-Point Source Contaminants Include Nitrate/Ammonia and Radionuclides



Summary

- **Historically, focus was on point source contamination, impact of poorly sealed wells easier to predict, easier to prevent.**
- **Many emerging contaminants are non-point source, by nature these are poorly constrained.**
- **Impact of poorly sealed wells difficult to predict and, thus very difficult to prevent.**